



ORIGINAL ARTICLE

Timing of elective pre-labour caesarean section: A decision analysis

Freke A. Wilmink¹ , Clarabelle T. Pham² , Nicole Edge³,
Chantal W.P.M. Hukkelhoven⁴, Eric A.P. Steegers⁵ and Ben W. Mol⁶

¹Department of Obstetrics and Gynaecology, Radboudumc, Nijmegen, The Netherlands

²School of Public Health, University of Adelaide, Adelaide, South Australia, Australia

³Department of Obstetrics and Gynaecology, Mildura Base Hospital, Mildura, Victoria, Australia

⁴Perined, Utrecht, The Netherlands

⁵Department of Obstetrics and Gynaecology, Erasmus MC – Sophia Children's Hospital, Rotterdam, The Netherlands

⁶Department of Obstetrics and Gynaecology, The Robinson Research Institute, University of Adelaide, North Adelaide, South Australia, Australia

Correspondence: Freke A. Wilmink, Gynaecologist, Department of Obstetrics and Gynaecology, Radboud University Medical Centre, no. 791, PO Box 9101, 6500 HB, Nijmegen, The Netherlands. Email: freke.wilmink@radboudumc.nl

Conflict of Interest: The authors report no conflicts of interest.

Received: 1 October 2017;

Accepted: 27 March 2018

Background: Since caesarean sections (CSs) before 39⁺⁰ weeks gestation are associated with higher rates of neonatal respiratory morbidity, it is recommended to delay elective CSs until 39⁺⁰ weeks. However, this bears the risk of earlier spontaneous labour resulting in unplanned CSs, which has workforce and resource implications, specifically in smaller obstetric units.

Aim: To assess, in a policy of elective CSs from 39⁺⁰ weeks onward, the number of unplanned CSs to prevent one neonate with respiratory complications, as compared to early elective CS.

Materials and Methods: We performed a decision analysis comparing early term elective CS at 37^{+0–6} or 38^{+0–6} weeks to elective prelabour CS, without strict medical indication, at 39^{+0–6} weeks, with earlier unplanned CS, in women with uncomplicated singleton pregnancies. We used literature data to calculate the number of unplanned CSs necessary to prevent one neonate with respiratory morbidity.

Results: Planning all elective CSs at 39^{+0–6} weeks required 10.9 unplanned CSs to prevent one neonate with respiratory morbidity, compared to planning all elective CSs at 38^{+0–6} weeks. Compared to planning all elective CSs at 37^{+0–6} weeks we needed to perform 3.3 unplanned CSs to prevent one neonate with respiratory morbidity.

Conclusion: In a policy of planning all elective pre-labour CSs from 39⁺⁰ weeks of gestation onward, between three and 11 unplanned CSs have to be performed to prevent one neonate with respiratory morbidity. Therefore, in our opinion, fear of early term labour and workforce disutility is no argument for scheduling elective CSs <39⁺⁰ weeks.

KEYWORDS

caesarean section, decision analysis, neonatal respiratory morbidity, respiratory distress syndrome, transient tachypnoea of the newborn

INTRODUCTION

Rates of caesarean sections (CSs) are increasing in Australia with 33.4% of women birthing by CS in 2015, compared to 28.5% in 2003.^{1,2} Of all birthing women in 2015, 21% underwent CS prior to labour.² This has significant health implications for infants, as elective CSs are associated with higher rates of neonatal respiratory morbidity compared to intended vaginal delivery.^{3–5} As this discrepancy reduces with increasing gestational age until 39⁺⁰ weeks of gestation, clinical guidelines in Australia and New Zealand, the UK and the USA recommend delaying elective pre-labour CS until 39⁺⁰ weeks gestation.^{5–13} Alternative recommendations are verification of lung maturity or to administer corticosteroids in advance for those CSs planned prior to 39⁺⁰ weeks.^{7,11,14} In 2012 still 51.6% of pre-labour CS occurred before 39⁺⁰ weeks in Australia.² This percentage seems even higher in private hospitals (66.8%).¹⁵ On one hand, delay of elective CSs until 39 weeks bears the risk of earlier spontaneous labour resulting in an unplanned CS. Given that in Australia more than 25% of women give birth in regional or remote areas, this may have resource and workforce implications as in these locations staff are often not on-site continuously and an unplanned CS may require initiation of 'call-backs'.¹ On the other hand, these locations are more likely to have limited access to neonatal facilities and thus reducing the potential for neonatal complications is crucially important. Use of a decision analysis tree is a quantitative approach to clinical problem solving that utilises local, national and international data to estimate the probabilities of a certain outcome. In this decision analysis we assessed, in a policy of elective CSs from 39⁺⁰ weeks onward, the number of unplanned CS that would be required to prevent one infant with respiratory complications.

MATERIALS AND METHODS

Definitions of CS

An elective (pre-labour) CS is defined as a planned CS without strict medical indication and performed before the start of labour. An unplanned CS is defined as a CS performed before the scheduled date after start of labour.

Decision tree model

Decision tree modelling enables the comparison of the outcomes of alternative clinical strategies in the absence of clinical trials.^{16,17} A decision tree with a three-week time horizon was constructed to model possible clinical strategies for timing of birth for women requiring elective CS (Fig. 1). The decision tree begins with three main branches stemming from a decision node, which represents the choice of a clinical strategy. A series of branches from each of the clinical strategies represents the different paths and outcomes for particular combinations of events. Each path on the decision tree has a probability of being taken and an outcome (neonate

with or without respiratory morbidity). The sum of the main outcome from the series of branches for each strategy was then calculated and compared. We assumed that elective CSs are not planned before 37⁺⁰ weeks of gestation. The three main strategies, *without* the administration of antenatal corticosteroids, were:

1. elective CS planned at 39^{+0–6} weeks of gestation, with risk of an unplanned CS at 37⁺⁰ until 38⁺⁶ weeks of gestation
2. elective CS booked at 38^{+0–6} weeks of gestation, with risk of an unplanned CS between 37^{+0–6} weeks of gestation
3. elective CS booked at 37^{+0–6} weeks of gestation.

Alternative strategies, *with* the administration of antenatal corticosteroids, were:

1. elective CS booked at 38^{+0–6} weeks of gestation, with risk of an unplanned CS at 37^{+0–6} weeks of gestation
2. elective CS booked at 37^{+0–6} weeks of gestation.

A second decision tree was constructed to include the possibility of lung maturity testing, using probabilities based on the sensitivity and specificity of the Lecithin-Sphingomyelin (L/S) ratio test (figure not shown).

An alternative strategy with performing an L/S ratio test was:

1. elective CS booked at 37^{+0–6} weeks of gestation after performing an L/S ratio test.

Our main outcome measure was the number of unplanned CSs needed to prevent one neonate with respiratory morbidity defined as respiratory distress syndrome (RDS) or transient tachypnoea of the newborn (TTN).

Data sources

The probabilities that were used as inputs for the decision tree model were based on previous literature and pregnancy outcome data from state-based and national government publications.^{2,18} To obtain relevant data we searched PubMed: 'Caesarean Section' [Mesh] in combination with the following free search terms: 'elective' or 'respiratory morbidity' or 'corticosteroids'. In addition, our search was filtered by 'English' and 'last 10 years'. The definitive search was performed on 9 July, 2017 and produced 1786 hits. After scanning titles, abstracts and cross-references we identified five cohort studies and one Cochrane review which presented data about neonatal respiratory morbidity following term elective CS stratified by week of gestation from 37⁺⁰ weeks of gestation onward (Table 1).^{5,8–10,19,20} The rate of unplanned CSs is based on the rate of singleton live births with spontaneous start of labour, with all women still having to deliver as the denominator, in 2012 until 2014. This was 3.75% between 37⁺⁰ and 37⁺⁶ weeks and 12.39% between 37⁺⁰ and 39⁺⁰ weeks of gestation, which corresponds with data from international literature.^{8,10,20}




FIGURE 1 Decision tree. CS, caesarean section; RDS, respiratory distress syndrome; TTN, transient tachypnoea of the newborn

TABLE 1 Neonatal respiratory morbidity associated with elective caesarean section

Reference	n	Respiratory outcome	Weeks		
			37 ⁺⁰⁻⁶ % (n/n _{total})	38 ⁺⁰⁻⁶ % (n/n _{total})	39 ⁺⁰⁻⁶ % (n/n _{total})
Morrison <i>et al.</i> ¹⁰	2341	NICU admission with RDS or TTN	7.4 (27/366)	4.2 (45/1063)	1.8 (9/505)
Hansen <i>et al.</i> ⁵	32 580	Combined respiratory morbidity†	10.0 (20/191)	5.1 (55/1083)	2.1 (22/1051)
Tita <i>et al.</i> ⁹	13 258	RDS or TTN	8.2 (68/833)	5.5 (213/3904)	3.5 (221/6510)
		RDS	3.7 (31/833)	1.9 (75/3904)	0.9 (58/6510)
		TTN	4.8 (40/833)	3.9 (153/3904)	2.7 (178/6510)
Wilmink <i>et al.</i> ⁸	20 973	Combined respiratory morbidity‡	6.8 (118/1734)	3.5 (356/10 139)	2.1 (136/6647)
		RDS or TTN	4.6 (80/1734)	2.7 (277/10 139)	1.5 (98/6647)
		RDS	0.5 (8/1734)	0.2 (21/10 139)	0.1 (7/6647)
		TTN	4.3 (74/1734)	2.5 (256/10 139)	1.4 (91/6647)
Chiossi <i>et al.</i> ¹⁹	23 749	RDS or TTN	9.1 (118/1296)	6.4 (294/4601)	4.0 (274/6941)
		RDS	3.8 (49/1296)	2.1 (98/4601)	1.0 (71/6941)
		TTN	5.3 (69/1296)	4.3 (196/4601)	2.9 (203/6941)

NICU, neonatal intensive care unit; RDS, respiratory distress syndrome; TTN, transient tachypnoea of the newborn.

†ICD-10 codes including any respiratory distress, TTN, or persistent pulmonary hypertension.

‡RDS, TTN, pneumothorax, oxygen supplementation, intermittent positive pressure ventilation or continuous positive airway pressure.

Analysis

Following the decision tree the number of unplanned CSs needed to be performed to prevent one neonate with respiratory morbidity was calculated, based on the obtained data. The risks of delivering earlier with an unplanned CS were taken into account. We assumed incidence rates of respiratory morbidity in neonates born after unplanned CS to be equivalent to the incidence rates of respiratory morbidity after elective CS.

Sensitivity analyses

One-way sensitivity analyses were performed to address the uncertainty and examine the impact of the probabilities, at 37⁺⁰⁻⁶, 38⁺⁰⁻⁶ and 39⁺⁰⁻⁶ weeks of gestation, of respiratory morbidity and unplanned CS on the cost to prevent one neonate with respiratory morbidity for our four main strategies. The lower and upper 95% limits were tested for the following parameters.

Probability of respiratory morbidity with:

- elective CS at 37⁺⁰⁻⁶ weeks *with* and *without* antenatal corticosteroids
- unplanned CS at 37⁺⁰⁻⁶ weeks *with* and *without* antenatal corticosteroids
- elective CS at 38⁺⁰⁻⁶ weeks *with* and *without* antenatal corticosteroids
- unplanned CS at 38⁺⁰⁻⁶ weeks *without* antenatal corticosteroids
- elective CS at 39⁺⁰⁻⁶ weeks *without* antenatal corticosteroids.

Probability of having an unplanned CS at:

- 37⁺⁰⁻⁶ weeks with a *planned elective* CS at 38⁺⁰⁻⁶ weeks
- 37⁺⁰⁻⁶ or 38⁺⁰⁻⁶ weeks with a *planned elective* CS at 39⁺⁰⁻⁶ weeks.

This study was exempt from institutional review board ethical approval.

RESULTS

Number of unplanned CSs needed to perform

Pooled incidences of RDS and TTN were calculated per week of gestation with separate outcome data from three large cohort studies.^{8,9,19} Pooled incidences of RDS and TTN together were 6.9%, 4.2% and 2.9% for neonates born at 37⁺⁰⁻⁶, 38⁺⁰⁻⁶ or 39⁺⁰⁻⁶ weeks of gestation, respectively (Table 2). This corresponds with a risk reduction of 39.1% by planning at 39⁺⁰⁻⁶ weeks compared to 38⁺⁰⁻⁶ weeks and of 58.0% by planning at 39⁺⁰⁻⁶ weeks compared to 37⁺⁰⁻⁶ weeks of gestation. We used data from the Cochrane review to calculate the risk reduction of respiratory morbidity after elective CS *with* the administration of antenatal corticosteroids. These relative risks (95% confidence intervals) were 0.49 (0.16-1.57) and 0.44 (0.17-1.14) at 37⁺⁰⁻⁶ and 38⁺⁰⁻⁶ weeks, respectively.²⁰

In Table 3 we present the results of our decision analysis.

Main strategies *without* the administration of antenatal corticosteroids:

1. when planning all elective CSs at 39⁺⁰⁻⁶ weeks, to prevent one neonate with respiratory morbidity we needed to perform:
 - a. 10.9 unplanned CSs, compared to planning all elective CSs at 38⁺⁰⁻⁶ weeks.

TABLE 2 Pooled incidences of RDS and TTN associated with elective caesarean section

Reference	n	Respiratory outcome	Weeks		
			37 ⁺⁰⁻⁶ %	38 ⁺⁰⁻⁶ %	39 ⁺⁰⁻⁶ %
Pooled incidences†	44 539	RDS or TTN	6.9	4.2	2.9
		RDS	2.3	1.0	0.7
		TTN	4.7	3.3	2.4

RDS, respiratory distress syndrome; TTN, transient tachypnea of the newborn.

†Pooled incidences with separate outcome data from Tita *et al.*,⁹ Wilmink *et al.*,⁸ and Choissi *et al.*¹⁹.

TABLE 3 Number of unplanned caesarean sections needed to be performed to prevent one neonate with RDS or TTN

Elective CS booked (weeks)	Number of emergency CS needed to perform†
Strategy <i>without</i> administration of antenatal corticosteroids	
For 39 ⁺⁰⁻⁶ compared to 38 ⁺⁰⁻⁶	10.9
For 39 ⁺⁰⁻⁶ compared to 37 ⁺⁰⁻⁶	3.3
For 38 ⁺⁰⁻⁶ compared to 37 ⁺⁰⁻⁶	1.4
Strategy <i>with</i> administration of antenatal corticosteroids	
For 38 ⁺⁰⁻⁶ <i>with</i> AC compared to 37 ⁺⁰⁻⁶ <i>with</i> AC	2.5
For 38 ⁺⁰⁻⁶ <i>with</i> AC compared to 39 ⁺⁰⁻⁶ <i>without</i> AC	3.0
Strategy <i>without</i> administration of antenatal corticosteroids after a positive L/S ratio	
For 39 ⁺⁰⁻⁶ compared to 37 ⁺⁰⁻⁶	3.9

AC, antenatal corticosteroids; CS, caesarean section; RDS, respiratory distress syndrome; TTN, transient tachypnoea of the newborn.

†Number of emergency CS needed to perform to prevent one neonate with RDS or TTN.

- b. 3.3 unplanned CSs, compared to planning all elective CSs at 37⁺⁰⁻⁶ weeks.
2. when planning all elective CSs at 38⁺⁰⁻⁶ weeks, to prevent one neonate with respiratory morbidity we needed to perform 1.4 unplanned CSs, compared to planning all elective CS at 37⁺⁰⁻⁶.

Strategies *with* the administration of antenatal corticosteroids:

1. when planning all elective CSs at 38⁺⁰⁻⁶ weeks *with* antenatal corticosteroids, to prevent one neonate with respiratory morbidity we needed to perform:
 - a. 2.5 unplanned CSs, compared to planning all elective CS at 37⁺⁰⁻⁶ *with* antenatal corticosteroids
 - b. 3.0 unplanned CSs, compared to planning all elective CS at 39⁺⁰ weeks *without* antenatal corticosteroids.

Strategy with performing an L/S ratio test:

1. when planning all elective CSs at 39 weeks, to prevent one neonate with respiratory morbidity we needed to perform 3.9 unplanned CSs compared to planning all elective CSs at 37⁺⁰⁻⁶ weeks after a positive L/S ratio test (and therefore assuming sufficient lung maturity).

DISCUSSION

Main findings

With a policy change of booking all elective CSs at 39⁺⁰⁻⁶ weeks of gestation, depending on prior booking policy at 37⁺⁰⁻⁶ or at 38⁺⁰⁻⁶ weeks of gestation, between three and 11 unplanned CSs are necessary to prevent one neonate with respiratory morbidity, respectively. Correspondingly, this will reduce the risk of a neonate with respiratory morbidity by approximately 50%. If delaying until 39⁺⁰ weeks is absolutely not possible because of fetal or maternal complications, a strategy with the administration of antenatal corticosteroids and booking an elective CS at 38⁺⁰⁻⁶ weeks compared to booking an elective CS at 37⁺⁰⁻⁶ weeks of gestation, 2.5 unplanned CSs need to be performed to prevent one neonate with respiratory morbidity. To prevent one neonate with respiratory morbidity, 3.9 unplanned CSs need to be performed if delaying until 39⁺⁰ weeks of gestation, compared to delivery at 37⁺⁰⁻⁶ weeks after a positive L/S ratio test (indicates fetal lung immaturity).

Strength and limitations

Our decision analysis shows valid and clear results and can assist in common practice and informed clinical decision-making. Large randomised trials are lacking and absolute numbers of severe respiratory morbidity at term are low. Therefore, we used data of large observational studies, published in peer-reviewed journals. All showed the same decreasing trend of respiratory morbidity with an increase of gestational age from 37⁺⁰ to 39⁺⁰ weeks onward, assuming validity. There were some uncertainties. We had no data on the incidence of respiratory morbidity in neonates born with unplanned CS after onset of labour; therefore we assumed these incidences to be equal, in agreement with recent literature.²¹ If this assumption is not correct (literature shows conflicting results),¹⁰ and neonatal respiratory

outcome after an intrapartum CS would be better compared to neonatal respiratory outcome after a prelabour CS, the number of CS needed to perform to prevent one sick neonate would be even lower. If the incidences of neonatal respiratory morbidity in CSs after the onset of labour were lower than after a planned elective CS, the numbers of unplanned CSs to perform to prevent one neonate with respiratory complications would be even lower. As numbers were too small to have valid data on the incidence of RDS and TTN, risk reduction of respiratory morbidity after administration of antenatal corticosteroids was calculated based on 'admission to a special baby care unit with respiratory morbidity'.²⁰ Although there are several adverse neonatal outcome measures associated with early term elective CSs (hypoglycaemia, hyperbilirubinaemia, sepsis, longer hospitalisation and neonatal intensive care admission),⁸ we only assessed respiratory morbidity as a neonatal outcome measure, as this is the most important cause of neonatal morbidity in early term births.

Interpretation

The number of unplanned CSs needing to perform will be translated into increased unplanned workforce and will have resource implications, but not necessarily only out of office hours. In a recent trial only 34% with a planning strategy at 39⁺³ and 36% with a planning strategy at 38⁺³ weeks of unplanned CSs was out of hours.²² One could argue that besides resource implications, unplanned CSs might have higher maternal risks compared with elective CSs. However, a large analysis did not show this.²³ In addition to respiratory morbidity, in CSs before 39⁺⁰ weeks, the risk for hypoglycaemia and longer neonatal admission is also significantly higher, causing maternal-neonatal separation which is associated with reduced rates of breast-feeding initiation.^{8,9,24} Also postnatal transfer to a more equipped neonatal unit can be necessary. In addition, a growing body of evidence suggests that children born early term have a greater risk for developmental delay in the first two years of life, have higher chances of respiratory morbidity like wheezing and asthma and more often have special educational needs.^{25,26,27–29}

Antenatal corticosteroids

A commonly used alternative strategy is administration of antenatal corticosteroids to all women with a planned CS before 39⁺⁰ weeks. A randomised clinical trial comparing all women undergoing a CS at term with and without prior administration of antenatal corticosteroids did show a decrease of admittance to special baby care units, (relative risk 0.46, 95% CI: 0.23–0.93); however, this decrease was not separately significant for TTN or RDS.^{14,20} Follow up in childhood did not show less asthma in the treated group and school assessment showed children in the treated group to be significantly more often in the lowest quartile of academic ability.³⁰ Furthermore, a recent trial showed an increase of neonatal hypoglycaemia in late preterm infants in the treated group (24.0% vs 15.0%) which has been associated with impaired neurologic outcome in childhood.^{31,32} As the

incidence of respiratory morbidity at term is low, following a strategy with administration of antenatal corticosteroids to all mothers, more than 95% of mothers and their neonates will be unnecessarily exposed to possible long-term risks.

Lung maturity tests as the L/S ratio (sensitivity 74.6% and specificity of 82.5%) and only performing a CS when the result indicates mature lungs as an alternative is questionable. Bates *et al.* reported that early-term born infants, despite adequate fetal lung maturity with L/S ratio test, still had a significant higher risk of TTN compared to infants born at 39–40 weeks gestation.³³ Quantus, a new non-invasive lung maturity test, unfortunately has no better test performances (sensitivity 62.1%, specificity 91.3%) between 34⁺⁰ and 38⁺⁶ weeks.³⁴

Future research

We recommend a large prospective cohort study planning all elective CSs at or beyond 39⁺⁰ weeks of gestation, to assess both neonatal and maternal outcome of elective and all necessary unplanned CSs in the short and long term.

CONCLUSION

In a policy of planning all elective pre-labour CSs from 39⁺⁰ weeks of gestation onwards, between three and 11 unplanned CSs have to be performed to prevent one neonate with respiratory morbidity. Therefore, in our opinion, fear of early-term labour and workforce disutility is no argument for scheduling elective CSs <39 weeks. Further justification might be provided by conducting a comprehensive economic evaluation. Administration of corticosteroids instead of postponing an elective CS until after 39⁺⁰ weeks is no good alternative until more data about long-term consequences are available.

ACKNOWLEDGEMENTS

BWM is supported by a NHMRC Practitioner Fellowship (GNT1 082548), and reports consultancy for ObsEva, Merck and Guerbet.

REFERENCES

1. Australian Institute of Health and Welfare. *Australia's mothers and babies 2015 - in brief*. Perinatal statistics series no. 33. Cat. no. PER 91. Canberra: AIHW, 2017.
2. Hilder L, Zhichao Z, Parker M *et al.* *Australia's mothers and babies 2012*. Perinatal statistics series no. 30. Cat. no. PER 69. Canberra: AIHW, 2014.
3. Levine EM, Ghai V, Barton JJ, Strom CM. Mode of delivery and risk of respiratory diseases in newborns. *Obstet Gynecol* 2001; **97**: 439–442.
4. Kolas T, Saugstad OD, Daltveit AK *et al.* Planned cesarean versus planned vaginal delivery at term: comparison of newborn infant outcomes. *Am J Obstet Gynecol* 2006; **195**: 1538–1543.

5. Hansen AK, Wisborg K, Uldbjerg N, Henriksen TB. Risk of respiratory morbidity in term infants delivered by elective caesarean section: cohort study. *BMJ* 2008; **336**: 85–87.
6. National Institute for Health and Care Excellence. *Caesarean section (CG132)* [Internet], 2012. [Accessed 12 January 2018.] Available from URL: <http://guidance.nice.org.uk/CG132>.
7. The Royal Australian and New Zealand College of Obstetricians and Gynaecologists. *C-Obs 23: Timing of Elective Caesarean Section at Term* [Internet], 2014. [Accessed 12 January 2018.] Available from URL: <http://www.ranzcog.edu.au/doc/timing-of-elective-cs-at-term.html>.
8. Wilmink FA, Hukkelhoven CW, Lunshof S et al. Neonatal outcome following elective cesarean section beyond 37 weeks of gestation: a 7-year retrospective analysis of a national registry. *Am J Obstet Gynecol* 2010; **202**(250): e1–e8.
9. Tita AT, Landon MB, Spong CY et al. Timing of elective repeat cesarean delivery at term and neonatal outcomes. *N Engl J Med* 2009; **360**: 111–120.
10. Morrison JJ, Rennie JM, Milton PJ. Neonatal respiratory morbidity and mode of delivery at term: influence of timing of elective caesarean section. *Br J Obstet Gynaecol* 1995; **102**: 101–106.
11. NIH State-of-the-Science Conference Statement on cesarean delivery on maternal request. *NIH Consens State Sci Statements* 2006; **23**: 1–29. Available at: <http://consensus.nih.gov/2006/cesarean-statement.pdf>. Retrieved January 12th, 2018.
12. American College of Obstetricians and Gynecologists. ACOG committee opinion no. 559: Cesarean delivery on maternal request. *Obstet Gynecol* 2013; **121**: 904–907.
13. Royal College of Obstetricians & Gynaecologists. *Birth After Previous Caesarean Birth. Greentop Guideline no. 45*. [Internet], 2015. [Accessed 12 January 2018.] Available from URL: https://www.rcog.org.uk/globalassets/documents/guidelines/gtg_45.pdf.
14. Stutchfield P, Whitaker R, Russell I. Antenatal betamethasone and incidence of neonatal respiratory distress after elective caesarean section: pragmatic randomised trial. *BMJ* 2005; **331**: 662.
15. Adams N, Gibbons KS, Tudehope D. Public-private differences in short-term neonatal outcomes following birth by prelabour caesarean section at early and full term. *Aust N Z J Obstet Gynaecol* 2017; **57**: 176–185.
16. Drummond MF, Jefferson TO. Guidelines for authors and peer reviewers of economic submissions to the BMJ. The BMJ Economic Evaluation Working Party. *BMJ* 1996; **313**: 275–283.
17. Sonnenberg FA, Beck JR. Markov models in medical decision making: a practical guide. *Med Decis Making* 1993; **13**: 322–338.
18. Perined. *Perinatal care in The Netherlands*, 2014. Utrecht: Perined.
19. Chiossi G, Lai Y, Landon MB et al. Timing of delivery and adverse outcomes in term singleton repeat cesarean deliveries. *Obstet Gynecol* 2013; **121**: 561–569.
20. Sotiriadis A, Makrydimas G, Papatheodorou S, Ioannidis JP. Corticosteroids for preventing neonatal respiratory morbidity after elective caesarean section at term. *Cochrane Database Syst Rev* 2009; **4**: CD006614.
21. Glavind J, Milidou I, Uldbjerg N et al. Neonatal morbidity after spontaneous labor onset prior to intended cesarean delivery at term: a cohort study. *Acta Obstet Gynecol Scand* 2017; **96**: 479–486.
22. Glavind J, Kindberg SF, Uldbjerg N et al. Elective caesarean section at 38 weeks versus 39 weeks: neonatal and maternal outcomes in a randomised controlled trial. *BJOG* 2013; **120**: 1123–1132.
23. Tita AT, Lai Y, Landon MB et al. Timing of elective repeat cesarean delivery at term and maternal perioperative outcomes. *Obstet Gynecol* 2011; **117**: 280–286.
24. Tavoulari EF, Benetou V, Vlastarakos PV et al. Factors affecting breast-feeding initiation in Greece: what is important? *Midwifery* 2015; **31**: 323–331.
25. Boyle EM, Poulsen G, Field DJ et al. Effects of gestational age at birth on health outcomes at 3 and 5 years of age: population based cohort study. *BMJ* 2012; **344**: e896.
26. MacKay DF, Smith GC, Dobbie R, Pell JP. Gestational age at delivery and special educational need: retrospective cohort study of 407,503 schoolchildren. *PLoS Med* 2010; **7**: e1000289.
27. Dueker G, Chen J, Cowling C, Haskin B. Early developmental outcomes predicted by gestational age from 35 to 41 weeks. *Early Hum Dev* 2016; **103**: 85–90.
28. Edwards MO, Kotecha SJ, Lowe J et al. Early-term birth is a risk factor for wheezing in childhood: a cross-sectional population study. *J Allergy Clin Immunol* 2015; **136**(581–7): e2.
29. Harju M, Keski-Nisula L, Georgiadis L et al. The burden of childhood asthma and late preterm and early term births. *J Pediatr* 2014; **164**(295–9): e1.
30. Stutchfield PR, Whitaker R, Gliddon AE et al. Behavioural, educational and respiratory outcomes of antenatal betamethasone for term caesarean section (ASTECS trial). *Arch Dis Child Fetal Neonatal Ed* 2013; **98**: F195–F200.
31. Gyamfi-Bannerman C, Thom EA, Blackwell SC et al. Antenatal betamethasone for women at risk for late preterm delivery. *N Engl J Med* 2016; **374**: 1311–1320.
32. Montassir H, Maegaki Y, Ogura K et al. Associated factors in neonatal hypoglycemic brain injury. *Brain Dev* 2009; **31**: 649–656.
33. Bates E, Rouse DJ, Mann ML et al. Neonatal outcomes after demonstrated fetal lung maturity before 39 weeks of gestation. *Obstet Gynecol* 2010; **116**: 1288–1295.
34. Palacio M, Bonet-Carne E, Cobo T et al. Prediction of neonatal respiratory morbidity by quantitative ultrasound lung texture analysis: a multicenter study. *Am J Obstet Gynecol* 2017; **217**: 196.